


**AMENDMENTS TO THE CLAIMS:**

The following listing of claims replaces all prior listings, and all prior versions, of claims in the application:

**LISTING OF CLAIMS:**

1. (Previously presented) A method of fluorination which comprises fluorinating a monosaccharide or monosaccharide bonded to a base of a nucleic acid using a fluorinating agent represented by general formula (I):




wherein Y represents nitrogen atom, R<sup>1</sup> and R<sup>2</sup> represent hydrogen atom or an alkyl group having 1 to 4 carbon atoms, and R<sup>1</sup> and R<sup>2</sup> may be bonded to each other to form a ring, and R<sup>0</sup> represents a benzene ring or a pyridine ring, which has one or two substituents selected from the group consisting of alkyl group having 1 to 2 carbon atoms, halogen atom, methoxy group, amino group and nitro group, which may be the same as or different from each other.

2. (Original) A method of fluorination according to Claim 1, wherein, in general formula (I), Y represents nitrogen atom, R<sup>0</sup> represents 3-methoxyphenyl group or 2-methoxyphenyl group, and R<sup>1</sup> and R<sup>2</sup> represent ethyl group.

3. (Previously presented) A method of fluorination according to Claim 1, wherein the monosaccharide or monosaccharide bonded to a base of a nucleic acid is fluorinated by a thermal reaction.

4. (Currently amended) A method of fluorination which comprises fluorinating a monosaccharide or monosaccharide bonded to a base of a nucleic acid by bringing the monosaccharide or monosaccharide bonded to a base of a nucleic acid and a fluorinating agent represented by general formula (II):




wherein Y represents nitrogen atom or phosphorus atom, x represents hydrogen or halogen atom, R<sup>0</sup>, R<sup>1</sup> and R<sup>2</sup> represent hydrogen atom or an alkyl or aryl group which may have substituents, the atom and the groups represented by R<sup>0</sup>, R<sup>1</sup> and R<sup>2</sup> may be a same with or different from each other, and two or three of the groups represented by R<sup>0</sup>, R<sup>1</sup> and R<sup>2</sup> may be bonded to each other to form a ring into reaction with each other under irradiation with at least one of microwave and electromagnetic wave having a wavelength around a microwave region.

5. (Previously presented) A method of fluorination according to Claim 4, wherein the monosaccharide or monosaccharide bonded to a base of a nucleic acid is fluorinated by bringing the monosaccharide or monosaccharide bonded to a base of a nucleic acid and the fluorinating agent represented by general formula (I) into reaction with each other under irradiation with microwave having a frequency of 1 to 30 GHz.

6. (Cancelled).

7. (Currently amended) A method of fluorination according to Claim 4, wherein the fluorinating agent is a compound represented by general formula (III):



wherein R<sup>3</sup>, R<sup>4</sup>, and R<sup>5</sup>~~R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup>~~ each independently represent an alkyl or aryl group which may have substituents, X represents hydrogen atom or a halogen atom, and two or three of the groups represented by R<sup>3</sup>, R<sup>4</sup>, and R<sup>5</sup>~~R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup>~~ may be bonded to each other to form a cyclic structure.

8. (Original) A method of fluorination according to Claim 7, wherein, in general formula (III), R<sup>3</sup> represents an aryl group which may have substituents, X represents fluorine atom, and R<sup>4</sup> and R<sup>5</sup> represent an alkyl or aryl group having 1 to 32 carbon atoms which may have substituents.

9.-12. (Cancelled).

13. (Previously presented) A method of fluorination according to Claim 4, wherein the fluorinating agent is a compound represented by general formula (II) in which X represents fluorine atom.

14. (Currently amended) A method of fluorination according to Claim 4, wherein the fluorinating agent is a compound represented by general formula (II) in which X represents fluorine atom, Y represents nitrogen atom,  $R^0$  represents 3-methylphenyl group or 2-methoxyphenyl group, and  $R^1$  and  $R^2$   $R^1$  and  $R^2$  represent ethyl group.

15.-20. (Cancelled).

21. (Previously presented) A method of fluorination according to Claim 4, wherein the fluorination is conducted in a presence of an agent accelerating a reaction.

22.-24. (Cancelled).

25. (Previously presented) The method of fluorination according to Claim 1, wherein the monosaccharide is selected from the group consisting of glucose, fucose, N-acetylglucosamine, N-acetylgalactosamine, N-acetylneuraminic acid, erythrose, threose, ribose, arabinose, xylose, alose, lyxose, altrose, mannose, gulose, idose, galactose, talose, psicose, fructose, sorbose, tagatose, hexanose, apiose, and a deoxy sugar, an amino sugar, a thio sugar, a condensed sugar and an anhydride of the monosaccharide, and the monosaccharide bonded to a base of a nucleic acid is selected from the group consisting of a nucleoside, an oligonucleoside, ribonucleic acid and deoxyribonucleic acid.